


CIPROGRAM

CIPROFLOXACIN 250 MG, 500 MG, 750 MG TABLETS



The reference powerful
treatment for urinary tract infections

 **Avalon Pharma**



CIPROGRAM

CIPROFLOXACIN 250 MG, 500 MG, 750 MG TABLETS

The reference powerful treatment for urinary tract infections

Active against most recognized bacteria that cause gastroenteritis and urinary tract infection ⁷

Does not greatly affect the normal anaerobic bowel flora ⁷

Concentrates at high level within mucosa ⁷

Penetrates affect into macrophages ⁸

Dosage and Administration

Urinary tract infection: **250 to 500mg** for 7-14 days

Chronic bacterial prostatitis: **500mg** twice daily for 28 days

CIPROGRAM 3

CIPROFLOXACIN 250 MG, 500 MG, 750 MG TABLETS

1. What CIPROGRAM is and what it is used for:

CIPROGRAM is an antibiotic belonging to the fluoro quinolone family. The active substance is ciprofloxacin. Ciprofloxacin works by killing bacteria that cause infections. It only works with specific strains of bacteria.

2. Before you take CIPROGRAM:

Do not take CIPROGRAM if you are allergic (hypersensitive) to the active substance, to other quinolone drugs or to any of the other ingredients of CIPROGRAM (see section 6), taking tizanidine (see Section 2. Taking other medicines). Severe, sudden allergic reaction (an anaphylactic reaction/shock, angio-oedema). Pain and swelling in the joints. Diarrhoea may develop while you are taking anti-biotics, including CIPROGRAM.

3. How to take CIPROGRAM:

Your doctor will explain to you exactly how much CIPROGRAM you will have to take as well as how often and for how long.

This will depend on the type of infection you have and how bad it is. The treatment usually lasts from 5 to 21 days, but may take longer for severe infections.

Take the tablets exactly as your doctor has told you. Ask your doctor or pharmacist if you are not sure how many tablets to take and how to take CIPROGRAM.

4. Possible side effects:

Like all medicines, CIPROGRAM can cause side effects, although not everybody gets them. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, tell your doctor or pharmacist. Common side effects (between 1 and 10 in every 100 people are likely to get these): nausea, diarrhoea, joint pains in children.

5. How to store CIPROGRAM:

Keep out of the reach and sight of children. Store below 30°C.

6. Further information:

The active substance in CIPROGRAM is Ciprofloxacin. Each tablet contains 250/500/750 mg of ciprofloxacin hydrochloride equivalent to 250/500/750 mg ciprofloxacin.

The other ingredients are: Microcrystalline Cellulose, maize starch, sodium starch glycolate, colloidal silicon dioxide, talc Powder, magnesium stearate, HPMC 2910, titanium dioxide, macrogol.



7. Nathwani D, Wood MJ. ciprofloxacin and salmonella carriage, J Hosp Infect 1992;22:181-4.

8. Easmon CS, Crane JP, Blowers AE. Effect of ciprofloxacin on intracellular organisms: in-vitro and in-vivo studies. J Antimicrob Chemother 1986;18 (suppl D) 43-8.

9. CIPROGRAM summary of products characteristics, SPC/ revised on 12/2014.

10. Ciprogram patient information leaflet (PIL).

 **Avalon Pharma**

Middle East Pharmaceutical Industries Co. Ltd.
P.O. Box 301332 Riyadh 11372

Tel: +966 11 464 3924

Fax: +966 11 462 0471

Email: info@avalon.com.sa avalonpharmaceutical.com