

Preserve quality of life with a well-tolerated and cost-effective treatment option










Erectile Dysfunction (ED) and Lower Urinary Tract Symptoms (LUTS) associated with Benign Prostatic Hyperplasia (BPH) are conditions with both high prevalence and significant negative impact on patients' quality of life. The ability to treat both LUTS and ED together with one medication and lifestyle advice is worthy of consideration(1)





Middle East Pharmaceutical Industries Co. Ltd. P.O. Box 301332 Rivadh 11372 Tel: +966 11 464 3924

Fax: +966 11 462 0471

Email: info@avalon.com.sa avalonpharmaceutical.com

¹⁻ Kirby M, Chapple C, Jackson G, et al. Erectile dysfunction and lower urinary tract symptoms: a consensus on the importance of co-diagnosis. Int J Clin Pract. 2013;67(7):606-618. doi:10.1111/jicp.12176
2- Hatzimouratidis K. A review of the use of tadalafil in the treatment of benign prostatic hyperplasia in men with and without erectile dysfunction. Ther Adv Urol. 2014;6(4):1355-147. doi:10.1177/1756287214531639
3- Carson CC, Rosenberg M, Kissel J, Wong DG. Tadalafil - a therapeutic option in the management of BPH-LUTS. Int J Clin Pract. 2014;68(1):94-103. doi:10.1111/j.jcp.12305
4- WANG, Y, BAO, Y, LIU, J, DUAN, L. and CUI, Y, (2018), Tadalafil 5 mg Once Daily Improves Lower Urinary Tract Symptoms and Erectile Dysfunction: A Systematic Review and Meta-analysis. Lower Urinary Tract Symptoms, 10: 84-92. doi:10.1111/j.ts.12144
5- Mônica FZ, De Nucci G. Tadalafil for the treatment of benign prostatic hyperplasia. Expert Opin Pharmacother. 2019;20(8):929-937. doi:10.1080/14656566.2019.1589452