

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What **Sitaviva** is and what it is used for
2. What you need to know before you take **Sitaviva**
3. How to take **Sitaviva**
4. Possible side effects
5. How to store **Sitaviva**
6. Further information

1. What Sitaviva is and what it is used for

Sitaviva contains the active substance sitagliptin which is a member of a class of medicines called DPP-4 inhibitors (dipeptidyl peptidase-4 inhibitors) that lowers blood sugar levels in adult patients with type 2 diabetes mellitus.

This medicine helps to increase the levels of insulin produced after a meal and decreases the amount of sugar made by the body.

Your doctor has prescribed this medicine to help lower your blood sugar, which is too high because of your type 2 diabetes. This medicine can be used alone or in combination with certain other medicines (insulin, metformin, sulphonylureas, or glitazones) that lower blood sugar, which you may already be taking for your diabetes together with a food and exercise plan.

What is type 2 diabetes?
Type 2 diabetes is a condition in which your body does not make enough insulin, and the insulin that your body produces does not work as well as it should. Your body can also make too much sugar.

When this happens, sugar (glucose) builds up in the blood. This can lead to serious medical problems like heart disease, kidney disease, blindness, and amputation.

2. Before you take Sitaviva

Do not take Sitaviva
if you are allergic to sitagliptin or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions
Cases of inflammation of the pancreas (pancreatitis) have been reported in patients receiving sitagliptin (see section 4).

If you encounter blistering of the skin, it may be a sign for a condition called bullous pemphigoid.

Your doctor may ask you to stop **Sitaviva**.

Tell your doctor if you have or have had:

- a disease of the pancreas (such as pancreatitis)
- gallstones, alcohol dependence or very high levels of triglycerides (a form of fat) in your blood. These medical conditions can increase your chance of getting pancreatitis (see section 4).
- type 1 diabetes
- diabetic ketoacidosis (a complication of diabetes with high blood sugar, rapid weight loss, nausea or vomiting)
- any past or present kidney problems.
- an allergic reaction to **Sitaviva** (see section 4).

This medicine is unlikely to cause low blood sugar because it does not work when your blood sugar is low. However, when this medicine is used in combination with a sulphonylurea medicine or with insulin, low blood sugar (hypoglycaemia) can occur. Your doctor may reduce the dose of your sulphonylurea or insulin medicine.

Other medicines and Sitaviva
Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

In particular, tell your doctor if you are taking digoxin (a medicine used to treat irregular heart beat and other heart problems). The level of digoxin in your blood may need to be checked if taking with **Sitaviva**.

Children and adolescents
Children and adolescents below 18 years should not use this medicine. It is not known if this medicine is safe and effective when used in children and adolescents under 18 years of age.

Pregnancy and breastfeeding
If you are pregnant or breastfeeding, think you may be pregnant or planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

You should not take this medicine during pregnancy.

It is not known if this medicine passes into breast milk. You should not take this medicine if you are breast-feeding or plan to breast-feed.

Driving and using machines
This medicine has no or negligible influence on the ability to drive and use machines. However, dizziness and drowsiness have been reported, which may affect your ability to drive or use machines.

Taking this medicine in combination with medicines called sulphonylureas or with insulin can cause hypoglycaemia, which may affect your ability to drive and use machines or work without safe foothold.

3. How to take Sitaviva

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual recommended dose is:

- one 100 mg film-coated tablet
- once a day
- by mouth

If you have kidney problems, your doctor may prescribe lower doses (such as 25 mg or 50 mg).

You can take this medicine with or without food and drink.

Your doctor may prescribe this medicine alone or with certain other medicines that lower blood sugar.

Diet and exercise can help your body use its blood sugar better. It is important to stay on the diet and exercise recommended by your doctor while taking **Sitaviva**.

If you take more Sitaviva than you should
If you take more than the prescribed dosage of this medicine, contact your doctor immediately.

If you forget to take Sitaviva
If you miss a dose, take it as soon as you remember. If you do not remember until it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not take a double dose of this medicine.

If you stop taking Sitaviva
Continue to take this medicine as long as your doctor prescribes it so you can continue to help control your blood sugar. You should not stop taking this medicine without talking to your doctor first.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP taking **Sitaviva** and contact a doctor immediately if you notice any of the following serious side effects:

- Severe and persistent pain in the abdomen (stomach area) which might reach through to your back with or without nausea and vomiting, as these could be signs of an inflamed pancreas (pancreatitis).

If you have a serious allergic reaction (frequency not known), including rash, hives, blisters on the skin/peeling skin and swelling of the face, lips, tongue, and throat that may cause difficulty in breathing or swallowing, stop taking this medicine and call your doctor right away. Your doctor may prescribe a medicine to treat your allergic reaction and a different medicine for your diabetes.

Some patients have experienced the following side effects after adding sitagliptin to metformin:

Common (may affect up to 1 in 10 people): low blood sugar, nausea, flatulence, vomiting.

Uncommon (may affect up to 1 in 100 people): stomach ache, diarrhoea, constipation, drowsiness.

Some patients have experienced different types of stomach discomfort when starting the combination of sitagliptin and metformin together (frequency is common).

Some patients have experienced the following side effects while taking sitagliptin in combination with a sulphonylurea and metformin: Very common (may affect more than 1 in 10 people): low blood sugar.

Common: constipation.

Some patients have experienced the following side effects while taking sitagliptin and pioglitazone:

Common: flatulence, swelling of the hands or legs.

Some patients have experienced the following side effects while taking sitagliptin in combination with pioglitazone and metformin: Common: swelling of the hands or legs.

Some patients have experienced the following side effects while taking sitagliptin in combination with insulin (with or without metformin): Common: flu.

Uncommon: dry mouth.

Some patients have experienced the following side effects while taking sitagliptin alone in clinical studies, or during post-approval use alone and/or with other diabetes medicines:

Common: low blood sugar, headache, upper respiratory infection, stuffy or runny nose and sore throat, osteoarthritis, arm or leg pain.

Uncommon: dizziness, constipation, itching

Rare: reduced number of platelets

Frequency not known: kidney problems (sometimes requiring dialysis), vomiting, joint pain, muscle pain, back pain, interstitial lung disease, bullous pemphigoid (a type of skin blister).

5. How to store Sitaviva

- Keep this medicine out of the sight and reach of children.
 - Store below 30°C.
 - Do not use this medicine after the expiry date which is stated on the label and carton. The expiry date refers to the last day of that month.
 - Do not throw away any medicines via wastewater or household waste.
- Ask your pharmacist how to throw away medicines if you no longer use. These measures will help to protect the environment.

6. Further information

What Sitaviva contains
The active substance is sitagliptin.

Each film-coated tablet contains:

- Sitagliptin Phosphate Monohydrate, equivalent to 100 mg Sitagliptin.
- Sitagliptin Phosphate Monohydrate, equivalent to 50 mg Sitagliptin.

The other ingredients are: In the tablet core: Dibasic calcium phosphate Anhydrous, Microcrystalline Cellulose, Croscarmellose sodium, Sodium Stearyl fumarate, Magnesium stearate. The tablet film coating contains: Polyvinyl alcohol, Macrogol, Talc, Yellow iron oxide, Titanium dioxide, Red iron oxide.

What Sitaviva looks like and contents of the pack
Sitaviva 50 mg: Light beige color, round, biconvex film-coated tablet debossed with C3 on one side and plain on other side.
Sitaviva 100 mg: Brownish orange, round, biconvex, film-coated tablet plain on both sides.
Pack size: 30 film-coated tablets.

Marketing Authorisation Holder and Manufacturer:
Middle East Pharmaceutical Industries Co Ltd (Avalon Pharma)
P.O.Box 4180 Riyadh 11491, Kingdom of Saudi Arabia
2.D Industrial City, Riyadh, Kingdom of Saudi Arabia
Tel: +966 (11) 2653948 -2653427
Fax: +966 (11) 2654723

This leaflet was last approved in 12/2019

To report any side effect(s):
• **Saudi Arabia:**

The National Pharmacovigilance and Drug Safety Centre (NPC)
• Fax: +966-11-205-7662 • SFDA call center: 19999 • Toll free phone: 8002490000 • E-mail: npc.drug@sfdagov.sa • Website: https://sdc.sfga.gov.sa

Other GCC States:
Please contact the relevant competent authority.
THIS IS A MEDICAMENT
- Medicament is a product which affects your health and its consumption contrary to instructions is dangerous for you. - Follow strictly the doctor's prescription, the method of use and the instructions of the pharmacist who sold the medicament. - The doctor and the pharmacist are the experts in medicines, their benefits and risks. - Do not by yourself interrupt the period of treatment prescribed for you. - Do not repeat the same prescription without consulting your doctor. - Keep all medicaments out of reach of children.
Council of Arab Health Ministers, Union of Arab Pharmacists